
FIG.




်

Growth inhibition [%]

Growth inhibition [8]

contrast to Taxol, doxorubicin, vincristine or epothilone against the mouse leukemic subline L1210/VCR as against the normal cytotoxic activity same has the D-24851 uI • MDR

LL1210



Influence of D-24851 on the multidrug-resistant murine leukemia L1210 (dose 10% of the $\mathrm{LD}_{50})$

	Dose (mg/kg) L1210	L1210 ILS %	L1210/VCR ILS %
D-24851	4 x 100 p.o. 4 x 147 p.o.	46 94	42 85
Adriamycin	4 x 1 i.p.	158	9
Taxol	4 x 15 i.p.	82	9
Vincristine	4 x 0.2 i.p.	47	-11

FIG. 3





FIG. 4

FIG. 5

Incubation of the migration of MO4 cells by D-24851



• D-24851 inhibits the migration of MO4 cells in a dose-dependent manner From this, an antiinvasive and an antimetastatic action can be derived for D-24851.

Neurotoxicity

Vincristine Taxol 10x 0.2 mg/kg i.p. 10x 15 mg/kg i.p.	‡	‡	+ + +	= no effect
Vincristine 10x 0.2 mg/kg i.p.	+	+	++	++ p > 0.01 vs. control
D-24851 10x 20 mg/kg p.o		•		
	Ataxia (rat)	Traction (rat)	Reaction (rat)	$+$ p ≥ 0.05 VS. control

D-24851 shows no neurotoxity [sic] in maximally $p \ge 0.05 \text{ VS. control}$

antitumor-active doses in contrast to Taxol and

vincristine

FIG. 6





FIG. 7

FIG. 8

Angiogenesis in human endothelial cell culture Vital staining, 44 hours after induction of angiogenesis

DMSO control



0.1 μM D24851





FIG. 9

Angiogenesis in human endothelial cell culture Lethal staining, 22 hours after induction of angiogenesis DMSO control



0.1 μM D24851

